VOL. 697, NOS. 1 + 2 JOURNAL OF CHROMATOGRAPHY A 21 APRIL 1995

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

20TH INTERNATIONAL SYMPOSIUM ON CHROMATOGRAPHY, BOURNEMOUTH, 19–24 JUNE 1994

Foreword
by M.B. Evans (Hatfield, UK) and A.F. Fell (Bradford, UK) 1

LIQUID CHROMATOGRAPHY

Theoretical and general

Criteria for developing rugged high-performance liquid chromatographic methods
by P.F. Vanbel and B.L. Tilquin, (Brussels, Belgium) and P.J. Schoenmakers (Amsterdam, Netherlands) 3

Systematic method development in hydrophobic interaction chromatography. I. Characterization of the phase system and modelling retention
by G. Rippel, A. Bede and L. Szepesy (Budapest, Hungary) 17

Expert system for the ion chromatographic determination of alkali and alkaline earth metals in mineral waters
by N. Gros and B. Gorenc (Ljubljana, Slovenia) 31

Performance of reversed-phase parallel-current open-tubular liquid chromatography columns and comparison with theory
by M. Horká, V. Kahle, M. Krejčí and K. Šlais (Brno, Czech Republic) 45

Effect of some mobile phase additives on the retention characteristics of different solute types on reversed-phase media. II
by S.D. McCrossen (Tonbridge, UK) and C.F. Simpson (London, UK) 53

Effect of various organic modifiers on the determination of the hydrophobicity parameters of non-homologous series of anticancer drugs
by E. Forgács and T. Cserháti (Budapest, Hungary) 59

Prediction of retention for substituted and unsubstituted polycyclic aromatic hydrocarbons in micellar liquid chromatography in the presence of organic modifiers
by M.A. Rodríguez Delgado, M.J. Sánchez, V. González and F. García Montelongo (La Laguna, Spain) 71

Evaluation of different techniques for peak purity assessment on a diode-array detector in liquid chromatography
by H. Fabre, A. Le Bris and M.D. Blanchin (Montpellier, France) 81

Sample preparation

Retention characteristics of octadecylsiloxane-bonded silica and porous polymer particle-loaded membranes for solid-phase extraction
by M.L. Mayer, S.K. Poole and C.F. Poole (Detroit, MI, USA) 89

Development of a binary solid-phase extraction cartridge for use in screening water samples for organic pollutants
by M.W. Powell (Pineham, UK) 101

Studies on trapping efficiencies of various collection devices for off-line supercritical fluid extraction
by N. Hüsers and W. Kleiböhmer (Münster, Germany) 107

High-performance liquid chromatography comparison of supercritical-fluid extraction and solvent extraction of microbial fermentation products
by S. Cocks, S.K. Wrigley and M.I. Chicarelli-Robinson (Slough, UK) and R.M. Smith (Loughborough, UK) . . . 115

Solid-phase extraction of polycyclic aromatic hydrocarbons from soil samples
by P.R. Kootstra, M.H.C. Straub, G.H. Stil and E.G. van der Velde (Bilthoven, Netherlands) and W. Hesselink and C.C.J. Land (Deventer, Netherlands) 123

Development and optimisation of an immunoaffinity-based solid-phase extraction for chlortoluron
by S.J. Shahtaheri, M.F. Katmeh, P. Kwasowski and D. Stevenson (Guildford, UK) 131

Trace enrichment by solid-phase extraction for the analysis of heavy metals in water
by V. Leepipatpiboon (Bangkok, Thailand) 137

Simplified clean-up for the determination of benzimidazolic fungicides by high-performance liquid chromatography with UV detection
by A. Di Muccio, I. Camoni, M. Ventriglia, D.A. Barbini, M. Mauro, P. Pelosi, T. Generali, A. Ausili and S. Girolimetti (Rome, Italy) 145

Columns

Effect of calcium-modified silica on retention and selectivity in normal-phase liquid chromatography
by M. Okamoto (Gifu, Japan), K. Nobuhara (Aichi, Japan) and D. Ishii (Kumamoto, Japan) 153

Stability of high-performance liquid chromatography columns packed with C_1 and C_8 polysiloxanes sorbed into porous silica particles
by T.A. Anazawa, F. Carraro, K.E. Collins and I.C.S.F. Jardim (Campinas, Brazil) 159

Separation of polycyclic aromatic hydrocarbons on a wide-pore polymeric C_{18} bonded phase
by W. Hesselink and R.H.N.A. Schiffer (Deventer, Netherlands) and P.R. Kootstra (Bilthoven, Netherlands) 165

Separation of metronidazole, its major metabolites and their conjugates using dynamically modified silica
by U.G. Thomsen, C. Cornett, J. Tjørnelund and S.H. Hansen (Copenhagen, Denmark) 175

Qualitative, semi-quantitative and spectrophotometric determination of ruthenium(III) by solid-phase extraction with 3-hydroxy-2-methyl-1,4-naphthoquinone-4-oxime-loaded polyurethane foam columns
by M.S. El-Shahawi and M. Almehdi (Al-Ain, United Arab Emirates) 185

Application of a comparative evaluation of several reversed-phase columns to the automated analysis of candidate pharmaceuticals
by I.M. Mutton (Greenford, UK) 191

Automated solid-phase extraction and coupled-column reversed-phase liquid chromatography for the trace-level determination of low-molecular-mass carbonyl compounds in air
by P.R. Kootstra and H.A. Herbold (Bilthoven, Netherlands) 203

Enantiomer and isomer separations

Porous graphitic carbon for the chromatographic separation of O-tetraacetyl-β-D-glucopyranosyl isothiocyanate-derivatised amino acid enantiomers
by W.C. Chan, R. Micklewright and D.A. Barrett (Nottingham, UK) 213

Chromatographic behaviour of positional isomers on porous graphitic carbon
by Q.H. Wan, P.N. Shaw, M.C. Davies and D.A. Barrett (Nottingham, UK) 219

Liquid chromatographic determination of amino acid enantiomers by derivatization with *o*-phthaldialdehyde and chiral thiols. Applications with reference to food science
by H. Brückner, M. Langer, M. Lüpke and T. Westhauser (Stuttgart, Germany) and H. Godel (Waldbronn, Germany) 229

Simple stationary phases derived from gluconolactone for chiral high-performance liquid chromatography (Short communication)
by K.M. Maher and D.R. Taylor (Manchester, UK) and H.J. Ritchie (Runcorn, UK) 247

Enantiomeric separation by packed column chiral supercritical fluid chromatography
by J. Whatley (Welwyn Garden City, UK) 251

Rapid method development for the separation of enantiomers by means of chiral column switching
by J.A. Whatley (Welwyn Garden City, UK) 257

Chiral resolution of protein kinase C inhibitors by reversed-phase high-performance liquid chromatography on cellulose tris-3,5-dimethylphenylcarbamate
by J.A. Whatley (Welwyn Garden City, UK) 263

Chiral high-performance liquid chromatography with cellulose carbamate-coated phases. Influence of support surface chemistry on enantioselectivity
by S.J. Grieb and S.A. Matlin (Coventry, UK), A.M. Belenguer (Tonbridge, UK) and H.J. Ritchie (Runcorn, UK) 271

Preparation of silicas combined with optically active organic compounds: optical resolution of metal chelate complexes on the silica composites
by F. Mizukami (Ibaraki, Japan), H. Izutsu (Hyogo, Japan), T. Osaka, Y. Akiyama, N. Uiji, K. Moriya and K. Endo (Chiba, Japan), K. Maeda and Y. Kiyozumi (Ibaraki, Japan) and K. Sakaguchi (Chiba, Japan) 279

High-performance liquid chromatographic determination of the geometrical isomers of β-carotene in several foodstuffs
by C.R.L. Carvalho, P.R.N. Carvalho and C.H. Collins (Campinas, Brazil) 289

Other applications

Use of chromogenic and fluorescent oxycarbonyl chlorides as reagents for amino acid analysis by high-performance liquid chromatography
by H. Brückner and M. Lüpke (Stuttgart, Germany) 295

Determination of the cysteine derivatives N-acetylcysteine, S-carboxymethylcysteine and methylcysteine in pharmaceuticals by high-performance liquid chromatography
by F.Y. Tsai, C.J. Chen and C.S. Chien (Taipei, Taiwan) 309

Indirect detection of saccharides in reversed-phase liquid chromatography with highly alkaline mobile phases
by B. Lu, M. Stefansson and D. Westerlund (Uppsala, Sweden) 317

Soluble glycoproteins from sugar cane juice analysed by high-performance liquid chromatography and fluorescence emission
by M.E. Legaz and M.M. Pedrosa (Madrid, Spain), R. De Armas and M.M. Martinez (Havana, Cuba) and C. Vicente (Madrid, Spain) 329

High-performance frontal analysis for the study of protein binding of troglitazone (CS-045) in albumin solution and in human plasma
by A. Shibukawa, T. Sawada and C. Nakao (Kyoto, Japan), T. Izumi (Tokyo, Japan) and T. Nakagawa (Kyoto, Japan) 337

Determination of traces of herbicide mixtures in water by on-line solid-phase extraction followed by liquid chromatography with diode-array detection and multivariate self-modelling curve resolution
by S. Lacorte, D. Barceló and R. Tauler (Catalonia, Spain) 345

Simultaneous determination of 27 phenols and herbicides in water by high-performance liquid chromatography with multi-electrode electrochemical detection
by G. Achilli and G.P. Cellerino (Parabiago, Italy), G.M. d'Eril (Pavia, Italy) and S. Bird (St. Ives, UK) 357

Improved method for the determination of glyphosate in water
by M.P. Abdullah, J. Daud, K.S. Hong and C.H. Yew (Selangor, Malaysia) 363

Sensitive determination of the benzene metabolite S-phenylmercapturic acid in urine by high-performance liquid chromatography with fluorescence detection
by T. Einig and W. Dehnen (Düsseldorf, Germany) 371

Quantitative estimation of heterocyclic aromatic amines by ion-exchange chromatography and electrochemical detection
by M.M.C. Van Dyck, B. Rollmann and C. De Meester (Brussels, Belgium) 377

Use of solvent optimization software for rapid selection of conditions for reversed-phase high-performance liquid chromatography of nicotine and its metabolites
by S. Pichini, I. Altieri, A.R. Passa, M. Rosa, P. Zuccaro and R. Pacifici (Rome, Italy) 383

Quantitation of phenylbutazone and oxyphenbutazone in equine plasma by high-performance liquid chromatography with solid-phase extraction
by M.R. Taylor and S.A. Westwood (Newmarket, UK) 389

Development of an isocratic high-performance liquid chromatographic method for monitoring of ciprofloxacin photodegradation
by K. Torniainen (Turku, Finland) and E. Mäki (Helsinki, Finland) 397

Micro high-performance liquid chromatography for the determination of nicarbazin in chicken tissues, eggs, poultry feed and litter
by R. Draisci, L. Lucentini, P. Boria and C. Lucarelli (Rome, Italy) 407

GAS CHROMATOGRAPHY

Chemometric classification of the solvent properties (selectivity) of commonly used gas chromatographic stationary phases
by S.K. Poole and C.F. Poole (Detroit, MI, USA) 415

Application of principal component factor analysis to the cavity model of solvation to identify factors important in characterizing the solvent properties of gas chromatographic stationary phases
by S.K. Poole and C.F. Poole (Detroit, MI, USA) 429

New equation for specific retention volumes in capillary column gas chromatography
by R. Lebrón-Aguilar, J.E. Quintanilla-López, A.M. Tello, A. Fernández-Torres and J.A. García-Domínguez (Madrid, Spain) 441

Identification of furan fatty acids in human blood cells and plasma by multi-dimensional gas chromatography–mass spectrometry
by H.G. Wahl, A. Chrzanowski, C. Müller and H.M. Liebich (Tübingen, Germany) and A. Hoffmann (Mülheim an der Ruhr, Germany) 453

Simultaneous determination of glycerol and mono-, di- and triglycerides in vegetable oil methyl esters by capillary gas chromatography
by C. Plank and E. Lorbeer (Vienna, Austria) 461

Monitoring of lipase-catalyzed cleavage of acylglycerols by high-temperature gas chromatography
by T.L. Bereuter and E. Lorbeer (Vienna, Austria) 469

Enantiomer separation of α-campholene and fencholene derivatives by capillary gas chromatography on permethylated cyclodextrin phases. I. Compounds separable with single columns
by R. Reinhardt, A. Steinborn, W. Engewald, K. Anhalt and K. Schulze (Leipzig, Germany) 475

Enantiomer separation of α-campholene and fencholene derivatives by capillary gas chromatography on permethylated cyclodextrins. II. Compounds separable with coupled techniques
by A. Steinborn, R. Reinhardt, W. Engewald, K. Wyssuwa and K. Schulze (Leipzig, Germany) 485

Computerized capillary gas chromatographic identification and determination of Siberian fir oil constituents
by A. Orav, K. Kuningas and T. Kailas (Tallinn, Estonia) 495

Comparative study of Colombian citrus oils by high-resolution gas chromatography and gas chromatography–mass spectrometry
by C. Blanco Tirado, E.E. Stashenko, M.Y. Combariza and J.R. Martinez (Bucaramanga, Colombia) 501

Aerobic bacterial degradation of selected polyaromatic compounds and *n*-alkanes found in petroleum
by E. Šepič and H. Leskovšek (Ljubljana, Slovenia) and C. Trier (Drake Circus, UK) 515

Optimization of the gas stripping and cryogenic trapping method for capillary gas chromatographic analysis of traces of volatile halogenated compounds in drinking water
by D. Djozan and Y. Assadi (Tabriz, Iran) 525

MISCELLANEOUS SEPARATION METHODS

Chromatographic separation of continuous mixtures in distributed pores
by B.J. McCoy (Davis, CA, USA) 533

Freeze–thaw flow management: a novel concept for high-performance liquid chromatography, capillary electrophoresis, electrochromatography and associated techniques
by C.D. Bevan and I.M. Mutton (Greenford, UK) 541

Enantioselective capillary electrophoresis of amino acid derivatives on cyclodextrin. Evaluation of structure–resolution relationships
by W. Lindner and B. Böhs (Graz, Austria) and V. Seidel (Linz, Austria) 549

Optimization of enantiomeric separations in capillary electrophoresis by reversal of the migration order and using different derivatized cyclodextrins
by T. Schmitt and H. Engelhardt (Saarbrücken, Germany) 561

Analysis of the quaternary structure of catalase by capillary zone electrophoresis
by M.M. Pedrosa, A. Reyes, C. Vicente and M.E. Legaz (Madrid, Spain) 571

Preparative-scale supercritical fluid chromatography
by K.D. Bartle (Leeds, UK), C.D. Bevan (Greenford, UK), A.A. Clifford, S.A. Jafar and N. Malak (Leeds, UK) and M.S. Verrall (Betchworth, UK) 579

Enantiomer separations by supercritical fluid chromatography on a chiral stationary phase physically anchored to porous graphitic carbon (Short communication)
by S.M. Wilkins and D.R. Taylor (Manchester, UK) and R.J. Smith (Harlow, UK) 587

Temperature dependence of chiral discrimination in supercritical fluid chromatography and high-performance liquid chromatography
by R.J. Smith (Harlow, UK) and D.R. Taylor and S.M. Wilkins (Manchester, UK) 591

High-performance thin-layer chromatographic method to determine sorption of propofol to infusion containers
by H. Salomies and P. Lautala (Helsinki, Finland) and M. Toppila (Oulu, Finland) 597

AUTHOR INDEX 603